

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
19 September 2002 (19.09.2002)

PCT

(10) International Publication Number  
WO 02/072202 A1

(51) International Patent Classification: A61P 25/00, 25/24, A61K 31/53, C07D 403/00

(21) International Application Number: PCT/US02/06837

(22) International Filing Date: 6 March 2002 (06.03.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/275,403 13 March 2001 (13.03.2001) US

(71) Applicant: BRISTOL-MYERS SQUIBB PHARMA COMPANY [US/US]; Patent Department, P.O. Box 4000, Princeton, NJ 08543-4000 (US).

(72) Inventor: GILLIGAN, Paul, J.; 2629 Pennington Drive, Wilmington, DE 19810 (US).

(74) Agents: DOLAN, Peter, L. et al.; Patent Department, P.O. Box 4000, Princeton, NJ 08543-4000 (US).

(81) Designated States (*national*): AB, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 4-(2-BUTYLAMINO)-2,7-DIMETHYL-8-(2-METHYL-6-METHOXYPYRID-3-YL) PYRAZOLO-[1,5-A]-1,3,5-TRIAZINE, ITS ENANTIOMERS AND PHARMACEUTICALLY ACCEPTABLE SALTS AS CORTICOTROPIN RELEASING FACTOR RECEPTOR LIGANDS



(I)

(57) Abstract: Corticotropin releasing factor (CRF) antagonists of Formula (I), and its use in treating anxiety, depression, and other psychiatric, neurological disorders as well as treatment of immunological, cardiovascular or heart-related diseases and colonic hypersensitivity associated with psychopathological disturbance and stress.

WO 02/072202 A1

4-(2-BUTYLAMINO)-2,7-DIMETHYL-8-(2-METHYL-6-METHOXYPYRID-3-YL)PYRAZOLO-[1,5-A]-1,3,5-TRIAZINE, ITS ENANTIOMERS AND PHARMACEUTICALLY ACCEPTABLE SALTS AS CORTICOTROPIN RELEASING FACTOR RECEPTOR LIGANDS

## 5 Field of the Invention

This invention relates to a treatment of psychiatric disorders and neurological diseases including major depression, anxiety-related disorders, post-traumatic stress disorder, supranuclear palsy and  
10 feeding disorders as well as treatment of immunological, cardiovascular or heart-related diseases and colonic hypersensitivity associated with psycho-pathological disturbances and stress, by administration of 4-(2-Butylamino)-2,7-dimethyl-8-(2-methyl-6-methoxypyrid-3-  
15 yl)pyrazolo-[1,5-a]-1,3,5-triazine, its enantiomer and pharmaceutically acceptable salts as a corticotropin releasing factor receptor ligand.

## Background of the Invention

Corticotropin releasing factor (herein referred to  
20 as CRF), a 41 amino acid peptide, is the primary physiological regulator of proopiomelanocortin (POMC) - derived peptide secretion from the anterior pituitary gland [J. Rivier et al., Proc. Nat. Acad. Sci. (USA) 80:4851 (1983); W. Vale et al., Science 213:1394

(1981)]. In addition to its endocrine role at the pituitary gland, immunohistochemical localization of CRF has demonstrated that the hormone has a broad extrahypothalamic distribution in the central nervous system and produces a wide spectrum of autonomic, electrophysiological and behavioral effects consistent with a neurotransmitter or neuromodulator role in brain [W. Vale et al., Rec. Prog. Horm. Res. 39:245 (1983); G.F. Koob, Persp. Behav. Med. 2:39 (1985); E.B. De Souza et al., J. Neurosci. 5:3189 (1985)]. There is also evidence that CRF plays a significant role in integrating the response of the immune system to physiological, psychological, and immunological stressors [J.E. Blalock, Physiological Reviews 69:1 (1989); J.E. Morley, Life Sci. 41:527 (1987)].

Clinical data provide evidence that CRF has a role in psychiatric disorders and neurological diseases including depression, anxiety-related disorders and feeding disorders. A role for CRF has also been postulated in the etiology and pathophysiology of Alzheimer's disease, Parkinson's disease, Huntington's disease, progressive supranuclear palsy and amyotrophic lateral sclerosis as they relate to the dysfunction of CRF neurons in the central nervous system [for review see E.B. De Souza, Hosp. Practice 23:59 (1988)].

In affective disorder, or major depression, the concentration of CRF is significantly increased in the cerebrospinal fluid (CSF) of drug-free individuals [C.B. Nemeroff et al., Science 226:1342 (1984); C.M. Banki et al., Am. J. Psychiatry 144:873 (1987); R.D. France et al., Biol. Psychiatry 28:86 (1988); M. Arato et al., Biol Psychiatry 25:355 (1989)]. Furthermore, the density of CRF receptors is significantly decreased in the frontal cortex of suicide victims, consistent with a hypersecretion of CRF [C.B. Nemeroff et al., Arch. Gen. Psychiatry 45:577 (1988)]. In addition, there is a blunted adrenocorticotropin (ACTH) response to CRF (i.v. administered) observed in depressed patients [P.W. Gold et al., Am J. Psychiatry 141:619 (1984); F. Holsboer et al., Psychoneuroendocrinology 9:147 (1984); P.W. Gold et al., New Eng. J. Med. 314:1129 (1986)]. Preclinical studies in rats and non-human primates provide additional support for the hypothesis that hypersecretion of CRF may be involved in the symptoms seen in human depression [R.M. Sapolsky, Arch. Gen. Psychiatry 46:1047 (1989)]. There is preliminary evidence that tricyclic antidepressants can alter CRF levels and thus modulate the numbers of CRF receptors in

brain [Grigoriadis et al., Neuropsychopharmacology 2:53 (1989)].

It has also been postulated that CRF has a role in the etiology of anxiety-related disorders. CRF produces  
5 anxiogenic effects in animals and interactions between benzodiazepine / non-benzodiazepine anxiolytics and CRF have been demonstrated in a variety of behavioral anxiety models [D.R. Britton et al., Life Sci. 31:363 (1982); C.W. Berridge and A.J. Dunn Regul. Peptides  
10 16:83 (1986)]. Preliminary studies using the putative CRF receptor antagonist  $\alpha$ -helical ovine CRF (9-41) in a variety of behavioral paradigms demonstrate that the antagonist produces "anxiolytic-like" effects that are qualitatively similar to the benzodiazepines [C.W.  
15 Berridge and A.J. Dunn Horm. Behav. 21:393 (1987), Brain Research Reviews 15:71 (1990)].

Neurochemical, endocrine and receptor binding studies have all demonstrated interactions between CRF and benzodiazepine anxiolytics, providing further  
20 evidence for the involvement of CRF in these disorders. Chlordiazepoxide attenuates the "anxiogenic" effects of CRF in both the conflict test [K.T. Britton et al., Psychopharmacology 86:170 (1985); K.T. Britton et al., Psychopharmacology 94:306 (1988)] and in the acoustic

startle test [N.R. Swerdlow et al., Psychopharmacology 88:147 (1986)] in rats. The benzodiazepine receptor antagonist (Ro15-1788), which was without behavioral activity alone in the operant conflict test, reversed  
5 the effects of CRF in a dose-dependent manner while the benzodiazepine inverse agonist (FG7142) enhanced the actions of CRF [K.T. Britton et al., Psychopharmacology 94:306 (1988)].

The mechanisms and sites of action through which  
10 the standard anxiolytics and antidepressants produce their therapeutic effects remain to be elucidated. It has been hypothesized however, that they are involved in the suppression of the CRF hypersecretion that is observed in these disorders. Of particular interest is  
15 that preliminary studies examining the effects of a CRF receptor antagonist ( $\alpha$ -helical CRF9-41) in a variety of behavioral paradigms have demonstrated that the CRF antagonist produces "anxiolytic-like" effects qualitatively similar to the benzodiazepines [for review  
20 see G.F. Koob and K.T. Britton, In: Corticotropin-Releasing Factor: Basic and Clinical Studies of a Neuropeptide, E.B. De Souza and C.B. Nemeroff eds., CRC Press p221 (1990)].

It has been further postulated that CRF has a role  
25 in cardiovascular or heart-related diseases as well as

gastrointestinal disorders arising from stress such as hypertension, tachycardia and congestive heart failure, stroke, irritable bowel syndrome post-operative ileus and colonic hypersensitivity associated with  
5 psychopathological disturbance and stress [for reviews see E.D. DeSouza, C.B. Nemeroff, Editors; Corticotropin-Releasing Factor: Basic and Clinical Studies of a Neuropeptide, E.B. De Souza and C.B. Nemeroff eds., CRC Press p221 (1990) and C. Mailliot, M. Million, J.Y. Wei,  
10 A. Gauthier, Y. Tache, Gastroenterology, 119, 1569-1579 (2000)].

Over-expression or under-expression of CRF has been proposed as an underlying cause for several medical disorders. Such treatable disorders include, for  
15 example and without limitation: affective disorder, anxiety, depression, headache, irritable bowel syndrome, post-traumatic stress disorder, supranuclear palsy, immune suppression, Alzheimer's disease, gastrointestinal diseases, anorexia nervosa or other  
20 feeding disorder, drug addiction, drug or alcohol withdrawal symptoms, inflammatory diseases, cardiovascular or heart-related diseases, fertility problems, human immunodeficiency virus infections, hemorrhagic stress, obesity, infertility, head and  
25 spinal cord traumas, epilepsy, stroke, ulcers,



amyotrophic lateral sclerosis, hypoglycemia, hypertension, tachycardia and congestive heart failure, stroke, osteoporosis, premature birth, psychosocial dwarfism, stress-induced fever, ulcer, diarrhea, post-operative ileus and colonic hypersensitivity associated with psychopathological disturbance and stress [for reviews see J.R. McCarthy, S.C. Heinrichs and D.E. Grigoriadis, *Curr. Pharm. Res.*, 5, 289-315 (1999); P.J. Gilligan, D.W. Robertson and R. Zaczek, *J. Medicinal Chem.*, 43, 1641-1660 (2000), G. P. Chrousos, *Int. J. Obesity*, 24, Suppl. 2, S50-S55 (2000); E. Webster, D.J. Torpy, I.J. Elenkov, G.P. Chrousos, *Ann. N.Y. Acad. Sci.*, 840, 21-32 (1998); D.J. Newport and C.B. Nemeroff, *Curr. Opin. Neurobiology*, 10, 211-218 (2000); G. Mastorakos and I. Ilias, *Ann. N.Y. Acad. Sci.*, 900, 95-106 (2000); M.J. Owens and C.B. Nemeroff, *Expert Opin. Invest. Drugs*, 8, 1849-1858 (1999); G. F. Koob, *Ann. N.Y. Acad. Sci.*, 909, 170-185 (2000)].

The following publications each describe CRF antagonist compounds; however, none disclose the compounds provided herein: WO95/10506; WO99/51608; WO97/35539; WO99/01439; WO97/44308; WO97/35846; WO98/03510; WO99/11643; PCT/US99/18707; WO99/01454; and, WO00/01675.

Summary of the Invention

In accordance with one aspect, the present invention provides a novel compound, pharmaceutical compositions and methods which may be used in the treatment of affective disorder, anxiety, depression, irritable bowel syndrome, post-traumatic stress disorder, supranuclear palsy, immune suppression, Alzheimer's disease, gastrointestinal disease, anorexia nervosa or other feeding disorder, drug or alcohol withdrawal symptoms, drug addiction, inflammatory disorder, fertility problems, disorders, the treatment of which can be effected or facilitated by antagonizing CRF, including but not limited to disorders induced or facilitated by CRF, or a disorder selected from inflammatory disorders such as rheumatoid arthritis and osteoarthritis, pain, asthma, psoriasis and allergies; generalized anxiety disorder; panic, phobias, obsessive-compulsive disorder; post-traumatic stress disorder; sleep disorders induced by stress; pain perception such as fibromyalgia; mood disorders such as depression, including major depression, single episode depression, recurrent depression, child abuse induced depression, and postpartum depression; dysthemia; bipolar disorders; cyclothymia; fatigue syndrome; stress-induced headache; cancer, human immunodeficiency virus (HIV) infections;

neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease and Huntington's disease; gastrointestinal diseases such as ulcers, irritable bowel syndrome, Crohn's disease, spastic colon, diarrhea, and post operative ilius and colonic hypersensitivity associated by psychopathological disturbances or stress; eating disorders such as anorexia and bulimia nervosa; hemorrhagic stress; stress-induced psychotic episodes; euthyroid sick syndrome; syndrome of inappropriate antidiarrhetic hormone (ADH); obesity; infertility; head traumas; spinal cord trauma; ischemic neuronal damage (e.g., cerebral ischemia such as cerebral hippocampal ischemia); excitotoxic neuronal damage; epilepsy; cardiovascular and hear related disorders including hypertension, tachycardia and congestive heart failure; stroke; immune dysfunctions including stress induced immune dysfunctions (e.g., stress induced fevers, porcine stress syndrome, bovine shipping fever, equine paroxysmal fibrillation, and dysfunctions induced by confinement in chickens, sheering stress in sheep or human-animal interaction related stress in dogs); muscular spasms; urinary incontinence; senile dementia of the Alzheimer's type; multiinfarct dementia; amyotrophic lateral sclerosis; chemical dependencies and

addictions (e.g., dependencies on alcohol, cocaine, heroin, benzodiazepines, or other drugs); drug and alcohol withdrawal symptoms; osteoporosis; psychosocial dwarfism; and hypoglycemia in a mammal.

5       The present invention provides a novel compound that binds to corticotropin releasing factor receptors, thereby altering the anxiogenic effects of CRF secretion. The compound of the present invention is useful for the treatment of psychiatric disorders and  
10       neurological diseases, anxiety-related disorders, post-traumatic stress disorder, supranuclear palsy and feeding disorders as well as treatment of immunological, cardiovascular or heart-related diseases and colonic hypersensitivity associated with psychopathological  
15       disturbance and stress in a mammal.

          According to another aspect, the present invention provides a novel compound of Formula (I) (described below) which is useful as an antagonist of the corticotropin releasing factor. The compound of the  
20       present invention exhibits activity as a corticotropin releasing factor antagonist and appears to suppress CRF hypersecretion. The present invention also includes pharmaceutical compositions containing such a compound of Formula (I), and methods of using such a compound for  
25       the suppression of CRF hypersecretion, and/or for the treatment of anxiogenic disorders.


The use of competitive binding assays is considered particularly valuable for screening candidates for new drugs, e.g. to identify new CRF ligands or other compounds having even greater or more selective binding  
5 affinity for CRF receptors, which candidates would therefore be potentially useful as drugs. In the assay, one determines the ability of the candidate ligand to displace the labelled compound.

Therefore, another embodiment of the invention  
10 includes the use of a compound of the invention is a binding assay, wherein one or more of the compounds may be joined to a label, where the label can directly or indirectly provide a detectable signal. Various labels include radioisotopes, fluorescers, chemilumescers,  
15 specific binding molecules, particles, e.g. magnetic particles, and the like.

Another embodiment of the invention is directed to the use of the compounds of the invention (particularly labeled compounds of this invention) as probes for the  
20 localization of receptors in cells and tissues and as standards and reagents for use in determining the receptor-binding characteristics of test compounds. Labeled compounds of the invention may be used for in vitro studies such as autoradiography of tissue sections  
25 or for in vivo methods, e.g. PET or SPECT scanning. Particularly, preferred compounds of the invention are useful as standards and reagents in determining the ability of a potential pharmaceutical to bind to the CRF1 receptor.

### Detailed Description of the Invention

[1] In a first embodiment, the present invention provides a compound of Formula (I):



(I)

and stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, and pharmaceutically acceptable salt or pro-drug forms thereof.

10 [2] In another embodiment, the present invention provides a compound of embodiment [1], isomers thereof, stereoisomeric forms thereof, mixtures of stereoisomeric forms thereof, pharmaceutically acceptable prodrugs thereof, or pharmaceutically acceptable salt forms thereof, wherein  
15 said compound is 4-((R)-2-butylamino)-2,7-dimethyl-8-(2-methyl-6-methoxypyrid-3-yl)-[1,5-a]-pyrazolo-1,3,5-triazine or 4-((S)-2-butylamino)-2,7-dimethyl-8-(2-methyl-6-methoxypyrid-3-yl)-[1,5-a]-pyrazolo-1,3,5-triazine.

[3] In another embodiment, the present invention  
20 provides a compound of any one of embodiments [1] to [2], pharmaceutically acceptable prodrugs thereof, or pharmaceutically acceptable salt forms thereof, wherein

said compound is substantially free of its (S) stereoisomer

[4] In another embodiment, the present invention provides a compound of embodiment [1], wherein said  
5 compound is 4-(2-butylamino)2,7-dimethyl-8-(2-methyl-6-methoxypyrid-3-yl)[1,5-a]-pyrazolo-1,3,5-triazine.

[5] In another embodiment, the present invention provides a compound of embodiment [1], wherein said compound is 4-((R)-2-butylamino)2,7-dimethyl-8-(2-  
10 methyl-6-methoxypyrid-3-yl)[1,5-a]-pyrazolo-1,3,5-triazine.

[6] A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of any  
15 one of embodiments [1] to [5].

[7] In another embodiment, the present invention provides a method of antagonizing a CRF receptor in a mammal, comprising administering to the mammal, a therapeutically effective amount of a compound of any  
20 one of embodiments [1] to [5].

[8] In another embodiment, the present invention provides a method of treating a disorder manifesting hypersecretion of CRF in a warm-blooded animal, comprising administering to the animal a therapeutically  
25 effective amount of a compound of any one of embodiments [1] to [5].

[9] In another embodiment, the present invention provides a method for the treatment of a disorder, the treatment of which can be effected or facilitated by  
30 antagonizing CRF, comprising administering to the mammal

a therapeutically effective of a compound of any one of embodiments [1] to [5].

[10] In another embodiment, the present invention provides a method of antagonizing a CRF receptor in a mammal, comprising administering to the mammal, a therapeutically effective amount of a compound of any one of embodiments [1] to [5].

[11] In another embodiment, the present invention provides a method of treating anxiety or depression in mammals, comprising administering to the mammal a therapeutically effective amount of a compound of any one of embodiments [1] to [5].

[12] In another embodiment, the present invention provides a method for screening for ligands for CRF receptors, which method comprises:

a) carrying out a competitive binding assay with a CRF receptor, a compound of any one of embodiments [1] to [5] which is labelled with a detectable label, and a candidate ligand; and

b) determining the ability of said candidate ligand to displace said labelled compound.

[13] In another embodiment, the present invention provides a method for detecting CRF receptors in tissue comprising:

a) contacting a compound of any one of embodiments [1] to [5], which is labelled with a detectable label, with a tissue, under conditions that permit binding of the compound to the tissue; and



b) detecting the labelled compound bound to the tissue.

[14] In another embodiment, the present invention provides a method of inhibiting the binding of CRF to a CRF-1 receptor, comprising contacting a compound of any one of embodiments [1] to [5] with a solution comprising cells expressing the CRF1 receptor, wherein the compound is present in the solution at a concentration sufficient to inhibit the binding of CRF to the CRF-1 receptor.

[15] In another embodiment, the present invention provides a article of manufacture comprising:

- a) a packaging material;
- b) a compound of any one of embodiments [1] to [5]; and
- c) a label or package insert contained within said packaging material idicating that said compound is effective for treating anxiety or depression.

[16] The present invention also comprises a method of treating affective disorder, anxiety, depression, headache, irritable bowel syndrome, post-traumatic stress disorder, supranuclear palsy, immune suppression, Alzheimer's disease, gastrointestinal diseases, anorexia nervosa or other feeding disorder, drug addiction, drug or alcohol withdrawal symptoms, inflammatory diseases, cardiovascular or heart-related diseases, fertility problems, human immunodeficiency virus infections, hemorrhagic stress, obesity, infertility, head and spinal cord traumas, epilepsy, stroke, ulcers,

amyotrophic lateral sclerosis, hypoglycemia or a disorder the treatment of which can be effected or facilitated by antagonizing CRF, including but not limited to disorders induced or facilitated by CRF, in  
5 mammals comprising administering to the mammal a therapeutically effective amount of a compound of any one of embodiments [1] to [5].

#### Definitions

As used herein, the term "pharmaceutically  
10 acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic acids, including inorganic acids and organic acids. Suitable non-toxic acids include inorganic and organic acids of basic residues such as amines, for example, acetic,  
15 benzenesulfonic, benzoic, amphorsulfonic, citric, ethenesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric  
20 acid, p-toluenesulfonic and the like; and alkali or organic salts of acidic residues such as carboxylic acids, for example, alkali and alkaline earth metal salts derived from the following bases: sodium hydride, sodium hydroxide, potassium hydroxide, calcium  
25 hydroxide, aluminum hydroxide, lithium hydroxide, magnesium hydroxide, zinc hydroxide, ammonia, trimethylammonia, triethylammonia, ethylenediamine, n-methyl-glucamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylenediamine, chloroprocaine,  
30 diethanolamine, procaine, n-benzylphenethylamine,

diethylamine, piperazine, tris(hydroxymethyl)-aminomethane, tetramethylammonium hydroxide, and the like.

Pharmaceutically acceptable salts of the compounds  
5 of the invention can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl  
10 acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in Remington's Pharmaceutical Sciences, 17th ed., Mack Publishing Company, Easton, PA, 1985, p. 1418, the disclosure of which is hereby incorporated by reference.

15 "Pharmaceutically acceptable prodrugs" as used herein means any covalently bonded carriers which release the active parent drug of Formula (I) in vivo when such prodrug is administered to a mammalian subject. Prodrugs of the compounds of Formula (I) are,  
20 within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals with undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended  
25 use, as well as the zwitterionic forms, where possible, of the compounds of the invention. The term "prodrug" means compounds that are rapidly transformed in vivo to

yield the parent compound of formula (I), for example by hydrolysis in blood. Functional groups which may be rapidly transformed, by metabolic cleavage, in vivo form a class of groups reactive with the carboxyl group of the compounds of this invention. They include, but are not limited to such groups as alkanoyl (such as acetyl, propionyl, butyryl, and the like), unsubstituted and substituted aroyl (such as benzoyl and substituted benzoyl), alkoxycarbonyl (such as ethoxycarbonyl), trialkylsilyl (such as trimethyl- and triethysilyl), monoesters formed with dicarboxylic acids (such as succinyl), and the like. Because of the ease with which the metabolically cleavable groups of the compounds useful according to this invention are cleaved in vivo, the compounds bearing such groups act as pro-drugs. The compounds bearing the metabolically cleavable groups have the advantage that they may exhibit improved bioavailability as a result of enhanced solubility and/or rate of absorption conferred upon the parent compound by virtue of the presence of the metabolically cleavable group. A thorough discussion of prodrugs is provided in the following: Design of Prodrugs, H. Bundgaard, ed., Elsevier, 1985; Methods in Enzymology, K. Widder et al, Ed., Academic Press, 42, p.309-396, 1985; A Textbook of Drug Design and Development, Krogsgaard-Larsen and H. Bundgaard, ed., Chapter 5; "Design and Applications of Prodrugs" p.113-191, 1991; Advanced Drug Delivery Reviews, H. Bundgard, 8, p.1-38, 1992; Journal of Pharmaceutical Sciences, 77, p. 285, 1988; Chem. Pharm. Bull., N. Nakeya et al, 32, p. 692,

1984; Pro-drugs as Novel Delivery Systems, T. Higuchi and V. Stella, Vol. 14 of the A.C.S. Symposium Series, and Bioreversible Carriers in Drug Design, Edward B. Roche, ed., American Pharmaceutical Association and  
5 Pergamon Press, 1987, which are incorporated herein by reference.

"Prodrugs" are considered to be any covalently bonded carriers which release the active parent drug of Formula (I) *in vivo* when such prodrug is administered to  
10 a mammalian subject. Prodrugs of the compounds of Formula (I) are prepared by modifying functional groups present in the compounds in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compounds.  
15 Prodrugs include compounds wherein hydroxy, amine, or sulfhydryl groups are bonded to any group that, when administered to a mammalian subject, cleaves to form a free hydroxyl, amino, or sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to,  
20 acetate, formate and benzoate derivatives of alcohol and amine functional groups in the compounds of Formula (I), and the like..

As used herein to describe a compound, the term "substantially free of its (S) stereoisomer" means that  
25 the compound is made up of a significantly greater proportion of its (R) stereoisomer than of its optical antipode (i.e., its (S) stereoisomer). In a preferred embodiment of the invention, the term "substantially free of its (S) stereoisomer" means that the compound is  
30 made up of at least about 90% by weight of its (R)

stereoisomer and about 10% by weight or less of its (S) stereoisomer.

In a more preferred embodiment of the invention, the term "substantially free of its (S) stereoisomer" means that the compound is made up of at least about 95% by weight of its (R) stereoisomer and about 5% by weight or less of its (S) stereoisomer. In an even more preferred embodiment, the term "substantially free of its (S) stereoisomer" means that the compound is made up of at least about 99% by weight of its (R) stereoisomer and about 1% or less of its (S) stereoisomer. In another preferred embodiment, the term "substantially free of its (S) stereoisomer" means that the compound is made up of nearly 100% by weight of its (R) stereoisomer. The above percentages are based on the total amount of the combined stereoisomers of the compound.

The term "therapeutically effective amount" of a compound of this invention means an amount effective to antagonize abnormal level of CRF or treat the symptoms of affective disorder, anxiety or depression in a host.

As used herein, the term "labeled" is meant that the compound is either directly or indirectly labeled with a label which provides a detectable signal, e.g. radioisotope, fluorescers, enzyme, antibodies, particles such as magnetic particles, chemiluminescer,  $P^{32}$ ,  $I^{131}$ , and  $At^{211}$ , etc.

### Syntheses

Many organic compounds exist in optically active forms, i.e., they have the ability to rotate the plane of plane-polarized light. In describing an optically  
5 active compound, the prefixes D and L or R and S are used to denote the absolute configuration of the molecule about its chiral center(s). The prefixes d and l or (+) and (-) are employed to designate the sign of rotation of plane-polarized light by the compound, with  
10 (-) or l meaning that the compound is levorotatory. A compound prefixed with (+) or d is dextrorotatory. For a given chemical structure, these compounds, called stereoisomers, are identical except that they are mirror images of one another. A specific stereoisomer may also  
15 be referred to as an enantiomer, and a mixture of such isomers is often called an enantiomeric mixture. A 50:50 mixture of enantiomers is referred to as a racemic mixture.


The present invention includes all stereoisomeric  
20 forms of the compounds of the formula I. Centers of asymmetry that are present in the compounds of formula I can all independently of one another have S configuration or R configuration. The prefixes d and l or (+) and (-) are employed to designate the sign of  
25 rotation of plane-polarized light by the compound, with (-) or l meaning that the compound is levorotatory. A compound prefixed with (+) or d is dextrorotatory. The invention includes all possible enantiomers and diastereomers and mixtures of two or more stereoisomers,  
30 for example mixtures of enantiomers and/or

diastereomers, in all ratios. Thus, enantiomers are a subject of the invention in enantiomerically pure form, both as levorotatory and as dextrorotatory antipodes, in the form of racemates and in the form of mixtures of the two enantiomers in all ratios. In the case of a cis/trans isomerism the invention includes both the cis form and the trans form as well as mixtures of these forms in all ratios. The preparation of individual stereoisomers can be carried out, if desired, by separation of a mixture by customary methods, for example by chromatography or crystallization, by the use of stereochemically uniform starting materials for the synthesis or by stereoselective synthesis. Optionally a derivatization can be carried out before a separation of stereoisomers. The separation of a mixture of stereoisomers can be carried out at the stage of the compounds of the formula I or at the stage of an intermediate during the synthesis. The present invention also includes all tautomeric forms of the compounds of formula (I).

The compound of Formula (I) may be prepared from using the procedures outlined in Scheme 1.

Scheme 1





A compound of Formula (II), where  $\text{X} = \text{F}$ , may be treated with a metal alkoxide (e.g. sodium methoxide, potassium methoxide; pre-formed or generated in situ) in an inert solvent to generate an intermediate of Formula (III). Inert solvents may include, but are not limited to, alkyl alcohols (1 to 8 carbons, preferably methanol or ethanol), lower alkanenitriles (1 to 6 carbons, preferably acetonitrile), water, dialkyl ethers (preferably diethyl ether), cyclic ethers (preferably

tetrahydrofuran or 1,4-dioxane), N,N-dialkylformamides (preferably dimethylformamide), N,N-dialkylacetamides (preferably dimethylacetamide), cyclic amides (preferably N-methylpyrrolidin-2-one), dialkylsulfoxides  
5 (preferably dimethylsulfoxide) or aromatic hydrocarbons (preferably benzene or toluene). Preferred reaction temperatures range from 0°C to 100°C.

Alternatively, a compound of Formula (II), where X = OH, may be treated with an alkylating agent in the  
10 presence of a base in an inert solvent to generate an intermediate of Formula (III). Alkylating agents include, but are not limited to, haloalkanes (e.g. CH<sub>3</sub>I), dialkyl sulfates (e.g. Me<sub>2</sub>SO<sub>4</sub>) or alkyl trifluoro-sulfonates (e.g. CH<sub>3</sub>O<sub>3</sub>SCF<sub>3</sub>).

15 Bases may include, but are not limited to, alkali metals, alkali metal hydrides (preferably sodium hydride), alkali metal alkoxides (1 to 6 carbons) (preferably sodium methoxide or sodium ethoxide), alkaline earth metal hydrides, alkali metal  
20 carbonates, alkaline metal carbonates, transition metal carbonates (e.g. silver carbonate), alkali metal dialkylamides (preferably lithium di-isopropylamide), alkali metal bicarbonates, alkali metal hydroxides, alkali metal bis(trialkylsilyl)amides (preferably sodium  
25 bis(trimethylsilyl)amide), trialkyl amines (preferably

N,N-di-isopropyl-N-ethyl amine) or aromatic amines (preferably pyridine).

Inert solvents may include, but are not limited to, halocarbons (1 to 8 carbons, 1 to 8 halogens), lower  
5 alkanenitriles (1 to 6 carbons, preferably acetonitrile), water, dialkyl ethers (preferably diethyl ether), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), N,N-dialkylformamides (preferably dimethylformamide), N,N-dialkylacetamides (preferably  
10 dimethylacetamide), cyclic amides (preferably N-methylpyrrolidin-2-one), dialkylsulfoxides (preferably dimethylsulfoxide) or aromatic hydrocarbons (preferably benzene or toluene). Preferred reaction temperatures range from 50°C to 150°C.

15 A compound of Formula (III) may be transformed to a compound of Formula (IV) by reaction with a brominating agent in the presence or absence of an additive in an inert solvent. Brominating agents include, but are not limited to, N-bromosuccinimide - 2,2'-azobisisobutyro-  
20 nitrile (AIBN), N-bromophthalimide - 2,2'-azobisisobutyronitrile (AIBN), bromine. Additives include, but are not limited to, alkali metal phosphates (e.g.  $K_3PO_4$ ,  $Na_3PO_4$ ), alkali metal hydrogen phosphates (e.g.  $Na_2HPO_4$ ,  $K_2HPO_4$ ), alkali metal dihydrogen phosphates (e.g.  
25  $NaH_2PO_4$ ,  $KH_2PO_4$ ). Inert solvents include, but are not

limited to, halocarbons (1 to 6 carbons, 1 to 6 halogens  
(preferably chlorine), water, N,N-dialkylformamides  
(preferably dimethylformamide), N,N-dialkylacetamides  
(preferably dimethylacetamide), cyclic amides  
5 (preferably N-methylpyrrolidin-2-one). Reaction  
temperatures range from 0°C to 200°C (preferably 20°C to  
120°C).

A compound of Formula (IV) may be converted to a  
compound of Formula (V) by sequential reactions with (1)  
10 an alkyl lithium in an inert solvent at temperatures  
ranging from -100°C to 50°C; (2) a compound of the  
Formula B(OR<sup>a</sup>), ( where R<sup>a</sup> is branched or straight chain  
alkyl of 1 to 20 carbons) at temperatures ranging from -  
100°C to 50°C and (3) an acid in the presence or absence  
15 of water at temperatures ranging from -100°C to 100°C.  
Alkyl lithiums may be branched or straight chain  
compounds containing 1 to 20 carbons. Inert solvents  
include, but are not limited to, dialkyl ethers  
(preferably diethyl ether), cyclic ethers (preferably  
20 tetrahydrofuran or 1,4-dioxane), or aromatic  
hydrocarbons (preferably benzene or toluene).

Acids may include, but are not limited to, alkanolic  
acids of 2 to 10 carbons (preferably acetic acid),  
haloalkanoic acids (2 - 10 carbons, 1-10 halogens, such  
25 as trifluoroacetic acid), arylsulfonic acids (preferably

p-toluenesulfonic acid or benzenesulfonic acid), alkanesulfonic acids of 1 to 10 carbons (preferably methanesulfonic acid), hydrochloric acid, sulfuric acid or phosphoric acid.

5 A compound of Formula (VII) may be produced by reaction of a compound of Formula (V) with a compound of Formula (VI) in the presence of a complex or salt of palladium or nickel, a base and an inert solvent. Complexes of palladium or nickel include, but are not  
10 limited to, phosphine complexes such as  $\text{Pd}(\text{PPh}_3)_4$ ,  $\text{PdCl}_2(\text{PPh}_3)_2$ ,  $\text{NiCl}_2(\text{PPh}_3)_2$ , or [1,1-bis(diphenylphosphino)ferrocene]-dichloropalladium. Bases may include, but are not limited to, alkali metals, alkali metal hydrides (preferably sodium  
15 hydride), alkali metal alkoxides (1 to 6 carbons) (preferably sodium methoxide or sodium ethoxide), alkali metal carbonates, alkaline metal carbonates (e.g. barium carbonate), transition metal carbonates (e.g. silver carbonate) or trialkyl amines  
20 (e.g. triethyl amine). Inert solvents may include, but are not limited to, dialkyl ethers (preferably diethyl ether), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), or aromatic hydrocarbons (preferably benzene or toluene). Preferred reaction temperatures  
25 range from  $-100^\circ\text{C}$  to  $100^\circ\text{C}$ .

An intermediate of Formula (VII) may be reacted with a base in the presence of an inert solvent to afford a compound of Formula (VIII), where M is an alkali metal cation (e.g. sodium or potassium). Bases  
5 may include, but are not limited to, alkali metal hydroxides (e.g. NaOH or KOH), alkali metal alkoxides (1 to 6 carbons) (preferably sodium methoxide or sodium ethoxide) or alkaline earth metal hydroxides. Inert solvents may include, but are not limited to, alkyl  
10 alcohols (1 to 6 carbons), lower alkanenitriles (1 to 6 carbons, preferably acetonitrile), water, cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), N,N-dialkylformamides (preferably dimethylformamide), N,N-dialkylacetamides (preferably dimethylacetamide), cyclic  
15 amides (preferably N-methylpyrrolidin-2-one), dialkylsulfoxides (preferably dimethylsulfoxide). Preferred reaction temperatures range from 0°C to 150°C.

Compounds of Formula (VIII) may be treated with hydrazine-hydrate in the presence of an acid and an  
20 inert solvent at temperatures ranging from 0°C to 200°C, preferably 70°C to 150°C, to produce compounds of Formula (IX). Acids may include, but are not limited to, alkanolic acids of 2 to 10 carbons (preferably acetic acid), haloalkanoic acids (2 - 10 carbons, 1-10  
25 halogens, such as trifluoroacetic acid), arylsulfonic

acids (preferably p-toluenesulfonic acid or benzenesulfonic acid), alkanesulfonic acids of 1 to 10 carbons (preferably methanesulfonic acid), hydrochloric acid, sulfuric acid or phosphoric acid.

5 Inert solvents may include, but are not limited to, water, alkyl alcohols (1 to 8 carbons, preferably methanol or ethanol), lower alkanenitriles (1 to 6 carbons, preferably acetonitrile), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), N,N-  
10 dialkylformamides (preferably dimethylformamide), N,N-dialkylacetamides (preferably dimethylacetamide), cyclic amides (preferably N-methylpyrrolidin-2-one), dialkylsulfoxides (preferably dimethylsulfoxide) or aromatic hydrocarbons (preferably benzene or toluene).

15 A compound of Formula (IX) may be reacted with compounds of Formula  $H_3C(C=NH)OR^C$  (where  $R^C$  is alkyl (1-6 carbons)) in the presence or absence of an acid in the presence of an inert solvent at temperatures ranging from 0°C to 200°C to produce a compound of Formula (X).

20 Acids may include, but are not limited to alkanoic acids of 2 to 10 carbons (preferably acetic acid), haloalkanoic acids (2 - 10 carbons, 1-10 halogens, such as trifluoroacetic acid), arylsulfonic acids (preferably p-toluenesulfonic acid or benzenesulfonic acid),  
25 alkanesulfonic acids of 1 to 10 carbons (preferably

methanesulfonic acid), hydrochloric acid, sulfuric acid or phosphoric acid. Stoichiometric or catalytic amounts of such acids may be used.

Inert solvents may include, but are not limited to, 5 water, alkanenitriles (1 to 6 carbons, preferably acetonitrile), halocarbons of 1 to 6 carbons and 1 to 6 halogens (preferably dichloroethane or chloroform), alkyl alcohols of 1 to 10 carbons (preferably ethanol), dialkyl ethers (4 to 12 carbons, preferably diethyl 10 ether or di-isopropylether) or cyclic ethers such as dioxan or tetrahydrofuran. Preferred temperatures range from 0°C to 100°C.

A compound of Formula (X) may be converted to an intermediate compound of Formula (XI) by treatment with 15 compounds  $C=O(R^d)_2$  (where  $R^d$  is halogen (preferably chlorine), alkoxy (1 to 4 carbons) or alkylthio (1 to 4 carbons)) in the presence or absence of a base in an inert solvent at reaction temperatures from -50°C to 200°C. Bases may include, but are not limited to, alkali 20 metal hydrides (preferably sodium hydride), alkali metal alkoxides (1 to 6 carbons) (preferably sodium methoxide or sodium ethoxide), alkali metal carbonates, alkali metal hydroxides, trialkyl amines (preferably N,N-di-



isopropyl-N-ethyl amine or triethylamine) or aromatic amines (preferably pyridine).

Inert solvents may include, but are not limited to, alkyl alcohols (1 to 8 carbons, preferably methanol or  
5 ethanol), lower alkanenitriles (1 to 6 carbons, preferably acetonitrile), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), N,N-dialkylformamides (preferably dimethylformamide), N,N-dialkylacetamides (preferably dimethylacetamide), cyclic amides  
10 (preferably N-methylpyrrolidin-2-one), dialkylsulfoxides (preferably dimethylsulfoxide) or aromatic hydrocarbons (preferably benzene or toluene).

A compound of Formula (XI) may be treated with a halogenating agent in the presence or absence of a base  
15 in the presence or absence of an inert solvent at reaction temperatures ranging from -80°C to 250°C to give a halogenated intermediate (XII) (where X is halogen). Halogenating agents include, but are not limited to, SOCl<sub>2</sub>, POCl<sub>3</sub>, PCl<sub>3</sub>, PCl<sub>5</sub>, POBr<sub>3</sub>, PBr<sub>3</sub> or  
20 PBr<sub>5</sub>. Bases may include, but are not limited to, trialkyl amines (preferably N,N-di-isopropyl-N-ethyl amine or triethylamine) or aromatic amines (preferably N,N-diethylaniline).

Inert solvents may include, but are not limited to,  
25 N,N-dialkylformamides (preferably dimethylformamide),

N,N-dialkylacetamides (preferably dimethylacetamide),  
cyclic amides (preferably N-methylpyrrolidin-2-one) or  
aromatic hydrocarbons (preferably benzene or toluene).  
Preferred reaction temperatures range from 20°C to  
5 200°C.

A compound of Formula (XII) may be reacted with an  
alkyl amine in the presence or absence of a base in the  
presence or absence of an inert solvent at reaction  
temperatures ranging from -80 to 250°C to generate  
10 compounds of Formula (I). Bases may include, but are not  
limited to, alkali metal hydrides (preferably sodium  
hydride), alkali metal alkoxides (1 to 6  
carbons) (preferably sodium methoxide or sodium  
ethoxide), alkaline earth metal hydrides, alkali metal  
15 dialkylamides (preferably lithium di-isopropylamide),  
alkali metal carbonates, alkali metal bicarbonates,  
alkali metal bis(trialkylsilyl)amides (preferably sodium  
bis(trimethylsilyl)amide), trialkyl amines (preferably  
N,N-di-isopropyl-N-ethyl amine) or aromatic amines  
20 (preferably pyridine).

Inert solvents may include, but are not limited to,  
alkyl alcohols (1 to 8 carbons, preferably methanol or  
ethanol), lower alkanenitriles (1 to 6 carbons,  
preferably acetonitrile), dialkyl ethers (preferably  
25 diethyl ether), cyclic ethers (preferably

tetrahydrofuran or 1,4-dioxane), N,N-dialkylformamides (preferably dimethylformamide), N,N-dialkylacetamides (preferably dimethylacetamide), cyclic amides (preferably N-methylpyrrolidin-2-one), dialkylsulfoxides  
5 (preferably dimethylsulfoxide), aromatic hydrocarbons (preferably benzene or toluene) or haloalkanes of 1 to 10 carbons and 1 to 10 halogens (preferably dichloroethane). Preferred reaction temperatures range from 0°C to 140°C.

10 The compounds of the invention may be prepared as radiolabeled compounds by carrying out their synthesis using precursors comprising at least one atom that is a radioisotope. The radioisotope is preferably selected from of at least one of carbon (preferably  $^{14}\text{C}$ ),  
15 hydrogen (preferably  $^3\text{H}$ ), sulfur (preferably  $^{35}\text{S}$ ), or iodine (preferably  $^{125}\text{I}$ ). Such radiolabeled probes are conveniently synthesized by a radioisotope supplier specializing in custom synthesis of radiolabeled probe compounds. Such suppliers include Amersham Corporation,  
20 Arlington Heights, Ill.; Cambridge Isotope Laboratories, Inc. Andover, Mass.; SRI International, Menlo Park, Calif.; Wizard Laboratories, West Sacramento, Calif.; ChemSyn Laboratories, Lexena, KS; American Radiolabeled Chemicals, Inc., St. Louis, Mo.; and Moravek Biochemicals  
25 Inc., Brea, Calif.

Tritium labeled probe compounds may also conveniently be prepared catalytically via platinum-catalyzed exchange in tritiated acetic acid, acid-

catalyzed exchange in tritiated trifluoroacetic acid, or heterogeneous-catalyzed exchange with tritium gas. Such preparations are also conveniently carried out as a custom radiolabeling by any of the suppliers listed in  
5 the preceding paragraph using the compound of the invention as substrate. In addition, certain precursors may be subjected to tritium-halogen exchange with tritium gas, tritium gas reduction of unsaturated bonds, or reduction using sodium borotritide, as appropriate.

10 Receptor autoradiography (receptor mapping) may be carried out in vitro as described by Kuhar in sections 8.1.1 to 8.1.9 of Current Protocols in Pharmacology (1998) John Wiley & Sons, New York, using radiolabeled compounds of the invention.

15 EXAMPLES

Analytical data were recorded for the compounds described below using the following general procedures. Proton NMR spectra were recorded on a Varian VXR or Unity 300 FT-NMR instruments (300 MHz); chemical shifts  
20 were recorded in ppm ( $\delta$ ) from an internal tetramethylsilane standard in deuteriochloroform or deuterodimethylsulfoxide as specified below. Mass spectra (MS) or high resolution mass spectra (HRMS) were recorded on a Finnegan MAT 8230 spectrometer or a  
25 Hewlett Packard 5988A model spectrometer (using chemi-ionization (CI) with  $\text{NH}_3$  as the carrier gas,

electrospray (ESI), atmospheric pressure chemi-  
ionization (APCI) or gas chromatography (GC)). Melting  
points were recorded on a MelTemp 3.0 heating block  
apparatus and are uncorrected. Boiling points are  
5 uncorrected. All pH determinations during workup were  
made with indicator paper.

Reagents were purchased from commercial sources  
and, where necessary, purified prior to use according to  
the general procedures outlined by D. Perrin and W.L.F.  
10 Armarego, Purification of Laboratory Chemicals, 3rd ed.,  
(New York: Pergamon Press, 1988). Chromatography was  
performed on silica gel using the solvent systems  
indicated below. For mixed solvent systems, the volume  
ratios are given. Otherwise, parts and percentages are  
15 by weight. Commonly used abbreviations are: DMF (N,N-  
dimethylformamide), EtOH (ethanol), MeOH (methanol),  
EtOAc (ethyl acetate), HOAc (acetic acid), DME (1,2-  
diethoxyethane) and THF (tetrahydrofuran).

The following examples are provided to describe the  
20 invention in further detail. These examples, which set  
forth the best mode presently contemplated for carrying  
out the invention, are intended to illustrate and not to  
limit the invention.

#### EXAMPLE 1

Preparation of 2,7-dimethyl-8-(2-methyl-6-methoxypyrid-3-yl)[1,5-a]-pyrazolo-[1,3,5]-triazin-4(3H)-one

A. 2-Methoxy-6-methylpyridine.

Sodium (31.0 g, 1.35 mol) was added portionwise to  
5 methanol (500 mL) over 30 min with stirring in a flask  
equipped with a reflux condenser. After the addition  
was complete, the reaction mixture was allowed to cool  
to ambient temperature. 2-Fluoro-6-methylpyridine (50  
g, 450 mmol) was added portionwise with stirring. The  
10 reaction mixture was then heated to reflux temperature  
and stirred for 48 h. The mix was then cooled to  
ambient temperature and solvent was removed in vacuo to  
provide a yellow oil. The residue was taken up in water  
(500 mL) and three extractions with ether (200 mL) were  
15 performed. The combined organic layers were dried over  
MgSO<sub>4</sub>, filtered and solvent was removed in vacuo from  
the filtrate to give a yellow liquid: <sup>1</sup>H-NMR(CDCl<sub>3</sub>, 300  
MHz): δ 7.44 (dd, 1H, J = 8, 7), 6.71 (d, 1H, J = 7),  
6.53 (d, 1H, J = 8), 3.91 (s, 3H), 2.45 (s, 3H).

20 B. 2-Methoxy-6-methylpyridine.

A mixture of 2-hydroxy-6-methylpyridine (6.85 g,  
62.8 mmol), silver carbonate (22.5 g, 81.6 mmol),  
iodomethane (39.1 mL, 628 mmol) and chloroform (200 mL)  
was stirred at ambient temperature for 40 h in the dark.  
25 The reaction mixture was filtered through celite. The

collected solid was washed with ether. The combined filtrates were concentrated in vacuo to give a liquid (6.25 g), which was identical to the product from Part A.

5 C. 6-Methoxy-3-bromo-2-methylpyridine.

A mixture of 2-methoxy-6-methylpyridine (17.0 g, 138 mmol) and a solution of disodium hydrogen phosphate (0.15M in water, 250 mL) was stirred at room temperature. Bromine (7.1 mL, 138 mmol) was added  
10 dropwise over 15 min via an addition funnel. The reaction mixture was then stirred at room temperature for 4 h. The clear colorless solution was diluted with water (500 mL) and extracted with dichloromethane (200 mL) three times. The combined organic layers were dried  
15 over  $\text{MgSO}_4$ , filtered and solvent was removed in vacuo from the filtrate to give a yellow liquid. Flash chromatography on silica gel (EtOAc:hexane::1:20) and removal of solvent from the desired combined fractions afforded a clear colorless liquid (15.4 g):  $^1\text{H-NMR}(\text{CDCl}_3$ ,  
20 300 MHz):  $\delta$  7.60(d, 1H,  $J = 8$ ), 6.46 (d, 1H,  $J = 8$ ), 3.89 (s, 3H), 2.54 (s, 3H).

D. 6-Methoxy-2-methylpyridine-3-boronic acid.

A solution of 6-methoxy-3-bromo-2-methylpyridine  
25 (59.8 g, 296 mmol) in dry THF (429 mL) was cooled with

stirring to ~ -78 °C under a nitrogen atmosphere. A solution of n-butyl lithium (2.5 M, 130.4 mL, 326 mmol) in hexane was added dropwise over 30 min. The reaction mixture was stirred for 3 h at ~ -78 °C. A solution of tri-isopropyl borate (102.7 mL, 445 mmol) in dry THF (100 mL) was added dropwise over 30 min. The reaction mixture was warmed to ambient temperature with stirring over 16 h. Acetic acid (37.35 g, 622 mmol), then water (110 mL) were added to the reaction mixture with stirring. After 2h, the layers were separated and the organic layer was concentrated in vacuo. The residue was taken up in 2-propanol (750 mL) and solvent was removed on a rotary evaporator (bath temperature ~ 50 °C). The residue was triturated with ether. The product was collected by filtration and dried in vacuo (48.4 g) : mp > 200 °C; <sup>1</sup>H-NMR(CD<sub>3</sub>OH, 300 MHz): δ 7.83 (d, 1H, J = 8), 6.56 (d, 1H, J = 8), 3.85 (s, 3H), 2.44 (s, 3H); GC-MS: 168 (M<sup>+</sup> + H).

20 E. 2-Methyl-3-(5-methylisoxazol-4-yl)-6-methoxypyridine.

A mixture of 4-iodo-5-methylisoxazole (18.2 g, 87 mmol), 6-methoxy-2-methylpyridine-3-boronic acid (14.6 g, 87 mmol), sodium bicarbonate (22.0 g, 262 mmol), water (150 mL) and DME (150 mL) was degassed three times



with stirring by the application of a vacuum and then introduction of a nitrogen atmosphere. [1,1-Bis(diphenylphosphino)ferrocene]-dichloropalladium (II) (2.14 g, 2.6 mmol) was added in one portion. The reaction mixture was degassed as before. The reaction mixture was then stirred at 80 °C for 4 h, then it was cooled to ambient temperature. Three extractions with EtOAc, drying the combined organic layers over MgSO<sub>4</sub>, filtration and removal of solvent in vacuo afforded an oil. Flash chromatography (EtOAc:hexane::1:9) and removal of solvent in vacuo from the desired fractions gave the product (7.15 g): <sup>1</sup>H-NMR(CDCl<sub>3</sub>, 300 MHz): δ 8.16 (s, 1H), 7.33 (d, 1H, J = 8), 6.63 (d, 1H, J = 8), 3.95 (s, 3H), 2.35 (s, 6H); APCI<sup>+</sup>-MS: 205 (M<sup>+</sup> + H).

15

F. 1-Cyano-1-(2-methyl-6-methoxypyrid-3-yl)propan-2-one, sodium salt.

A mixture of sodium methoxide (25% w/w, 13 mL, 70 mmol), 2-methyl-3-(5-methylisoxazol-4-yl)-6-methoxypyridine (7.15 g, 35 mmol) and methanol (50 mL) was stirred at room temperature for 16 h. Solvent was removed in vacuo to give a yellow oil. Trituration with ether, filtration and drying in vacuo afforded the crude product as a white solid (9.3 g).

25

G. 5-Amino-4-(2-methyl-6-methoxypyrid-3-yl)-3-methylpyrazole.

A mixture of 1-cyano-1-(2-methyl-6-methoxypyrid-3-yl)propan-2-one, sodium salt (9.3 g), hydrazine-hydrate  
5 (6 mL, 123.3 mmol) and glacial acetic acid (150 mL) was stirred at room temperature for 4 h. The reaction mixture was concentrated in vacuo. The residue was dissolved in 1N HCl and the resulting solution was extracted with EtOAc two times. A 1N NaOH solution was  
10 added to the aqueous layer until pH = 12. The resulting semi-solution was extracted three times with ethyl acetate. The combined organic layers were dried over MgSO<sub>4</sub> and filtered. Solvent was removed in vacuo to give a viscous oil (5.8 g): <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 300 MHz):  
15 7.37 (d, 2H, J = 8), 6.62 (d, 2H, J = 8), 3.95 (s, 3H), 2.36 (s, 3H), 2.08 (s, 3H); APCI<sup>+</sup>-MS: 219 (M<sup>+</sup> + H); 260 (M<sup>+</sup> + CH<sub>3</sub>CN).

H. 5-Acetamidino-4-(2-methyl-6-methoxypyrid-3-yl)-3-methylpyrazole, acetic acid salt.

Ethyl acetamidate hydrochloride (6.46g, 52.2 mmol) was added quickly to a rapidly stirred mixture of potassium carbonate (6.95g, 50.0 mol), dichloromethane (60 mL) and water (150 mL). The layers were separated  
25 and the aqueous layer was extracted with dichloromethane

(2 X 60 mL). The combined organic layers were dried over MgSO<sub>4</sub> and filtered. Solvent was removed by simple distillation and the pot residue, a clear pale yellow liquid, was used without further purification.

5        Glacial acetic acid (1.0 mL, 17.4 mmol) was added to a stirred mixture of 5-amino-4-(2-methyl-6-methoxypyrid-3-yl)-3-methylpyrazole (3.8g, 17.4 mmol), ethyl acetamidate free base and dichloromethane (100 mL). The resulting reaction mixture was stirred at room  
10    temperature for 16 h; at the end of which time, it was concentrated in vacuo. The residue was triturated with ether, the product was filtered and washed with copious amounts of ether. The white solid was dried in vacuo (5.4 g): <sup>1</sup>H-NMR (CD<sub>3</sub>OH, 300 MHz): 7.43 (d, 2H, J = 8),  
15    6.69 (d, 2H, J = 8), 4.9 (br s, 2H), 3.93 (s, 3H), 2.31 (s, 3H), 2.24 (s, 3H), 2.13 (s, 3H), 1.88 (s, 3H); APCI<sup>+</sup>-MS: 260 (M<sup>+</sup> + H).

I.    2,7-dimethyl-8-(2-methyl-6-methoxypyrid-3-yl)[1,5-a]-pyrazolo-[1,3,5]-triazin-4(3H)-one.

20        Sodium pellets (3.9 g, 169 mmol) were added portionwise to ethanol (200 mL) with vigorous stirring. After all the sodium reacted, 5-acetamidino-4-(2-methyl-6-methoxypyrid-3-yl)-3-methylpyrazole, acetic acid salt (5.4 g, 16.9 mmol) and diethyl carbonate ( 16.4 mL,  
25    135.3 mmol) were added. The resulting reaction mixture

was heated to reflux temperature and stirred for 18 hours. The mix was cooled to room temperature and solvent was removed in vacuo. The residue was dissolved in water and a 1N HCl solution was added slowly until pH 5 ~ 6. The aqueous layer was extracted with EtOAc three times; the combined organic layers were dried over MgSO<sub>4</sub> and filtered. Solvent was removed in vacuo to give a solid. Trituration with ether, filtration and drying in vacuo afforded a white solid (3.9 g): <sup>1</sup>H-NMR (CD<sub>3</sub>OH, 300 MHz): 7.49 (d, 2H, J = 8), 6.69 (d, 2H, J = 8), 3.93 (s, 3H), 2.35 (s, 3H), 2.28 (s, 3H), 2.24 (s, 3H); APCI<sup>+</sup>-MS: 286 (M<sup>+</sup> + H).

#### EXAMPLE 2

Preparation of 4-((R)-2-butylamino)2,7-dimethyl-8-(2-methyl-6-methoxypyrid-3-yl)[1,5-a]-pyrazolo-1,3,5-triazine

A. 4-Chloro-2,7-dimethyl-8-(2-methyl-6-methoxypyrid-3-yl)[1,5-a]-pyrazolotriazine.

A mixture of 2,7-dimethyl-8-(2-methyl-6-methoxypyrid-3-yl)[1,5-a]-pyrazolo-1,3,5-triazin-4-one (Example 1, 3.9 g, 13.7 mmol), di-isopropyl-ethylamine (9.5 mL, 54.7 mmol), phosphorus oxychloride (5.1 mL, 54.7 mmol) and toluene (75 mL) was stirred at reflux temperature for 4 h. The volatiles were removed in vacuo. The residue was loaded on a pad of silica gel on celite and eluted with a 1:1 mixture of EtOAc and

hexane. Solvent was removed in vacuo from the filtrate to give an oil.

B. 4-((R)-2-butylamino)2,7-dimethyl-8-(2-methyl-6-methoxypyrid-3-yl)[1,5-a]-pyrazolo-1,3,5-triazine.

5 A mixture of 4-chloro-2,7-dimethyl-8-(2-methyl-6-methoxypyrid-3-yl)[1,5-a]-pyrazolotriazine, (R)-2-butylamine (2.0 mL, 20.5 mmol), di-isopropyl-ethylamine (9.5 mL, 54.7 mmol) and dry THF (25 mL) was stirred at ambient temperature for 18 hours. Solvent was removed in  
10 vacuo. Column chromatography of the residue (first using EtOAc:hexane::1:2, then using EtOAc:hexane::1:4) afforded the product. Removal of solvent in vacuo gave a white solid (2.3 g): mp = 118.3 °C ; <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 300 MHz): δ 7.41 (d, 1H, J = 8), 6.63 (d, 1H, J = 8),  
15 6.25 (br d, 1H, J = 9), 4.35-4.30 (m, 1H), 3.95 (s, 3H), 2.49 (s, 3H), 2.35 (s, 3H), 2.30 (s, 3H), 1.76-1.66 (m, 2H), 1.34 (d, 3H, J = 7), 1.02 (t, 3H, J = 7); <sup>13</sup>C-NMR (CDCl<sub>3</sub>, 100.52 MHz): δ 163.8, 163.0, 155.7, 153.7, 147.8, 146.6, 141.6, 118.5, 107.4, 106.6, 53.3, 48.2,  
20 29.7, 26.1, 22.9, 20.4, 13.1, 10.3; IR (neat, KBr, cm<sup>-1</sup>): 3380 (m), 3371 (m), 2968 (m), 2928 (m), 2872 (w), 1621 (s), 1588 (s), 1544 (s), 1489 (s), 1460 (s), 1425 (s), 1413 (s), 1364 (s), 1346 (m), 1304 (s), 1275 (s), 1247 (s), 1198 (m), 1152 (m), 1134 (m), 1112 (m), 1034

(s), 1003 (m); ESI(+)-HRMS: Calcd for  $C_{18}H_{24}N_6O$ : 341.2089; Found: 341.2093 ( $M^+ + H$ ). Anal. Calcd for  $C_{18}H_{24}N_6O$ : C, 63.51, H, 7.12, N, 24.69; Found: C, 63.67, H, 7.00, N, 24.49.

5

### Utility

Rat CRF Receptor Binding Assay for the Evaluation of Biological Activity.

Receptor binding affinity to rat cortical receptors was assayed according to the published methods (E.B. De Souza, J. Neuroscience, 7: 88 (1987)).

Curves of the inhibition of [ $^{125}I$ -Tyr<sup>0</sup>]-o-CRF binding to cell membranes at various dilutions of test drug were analyzed by the iterative curve fitting program LIGAND [P.J. Munson and D. Rodbard, Anal. Biochem. 107:220 (1980)], which provides  $K_i$  values for inhibition which are then used to assess biological activity.

### Inhibition of CRF-Stimulated Adenylate Cyclase Activity

Inhibition of CRF-stimulated adenylate cyclase activity can be performed as described by G. Battaglia et al. Synapse 1:572 (1987). Briefly, assays are carried out at 37° C for 10 min in 200 ml of buffer containing 100 mM Tris-HCl (pH 7.4 at 37° C), 10 mM  $MgCl_2$ , 0.4 mM EGTA, 0.1% BSA, 1 mM

isobutylmethylxanthine (IBMX), 250 units/ml  
phosphocreatine kinase, 5 mM creatine phosphate, 100 mM  
guanosine 5'-triphosphate, 100 nM oCRF, antagonist  
peptides (concentration range  $10^{-9}$  to  $10^{-6}$ M) and 0.8 mg  
5 original wet weight tissue (approximately 40-60 mg  
protein). Reactions are initiated by the addition of 1  
mM ATP/ $^{32}$ P]ATP (approximately 2-4 mCi/tube) and  
terminated by the addition of 100 ml of 50 mM Tris-HCL,  
45 mM ATP and 2% sodium dodecyl sulfate. In order to  
10 monitor the recovery of cAMP, 1  $\mu$ l of [ $^3$ H]cAMP  
(approximately 40,000 dpm) is added to each tube prior  
to separation. The separation of [ $^{32}$ P]cAMP from  
[ $^{32}$ P]ATP is performed by sequential elution over Dowex  
and alumina columns.

#### 15 In vivo Biological Assay

The in vivo activity of a compound of the present  
invention can be assessed using any one of the  
biological assays available and accepted within the art.  
Illustrative of these tests include the Acoustic Startle  
20 Assay, the Stair Climbing Test, and the Chronic  
Administration Assay. These and other models useful for  
the testing of compounds of the present invention have  
been outlined in C.W. Berridge and A.J. Dunn Brain  
Research Reviews 15:71 (1990).

A compound may be tested in any species of rodent or small mammal.

A compound of this invention has utility in the treatment of imbalances associated with abnormal levels of corticotropin releasing factor in patients suffering from depression, affective disorders, and/or anxiety.

A compound of this invention can be administered to treat these abnormalities by means that produce contact of the active agent with the agent's site of action in the body of a mammal. The compounds can be administered by any conventional means available for use in conjunction with pharmaceuticals either as individual therapeutic agent or in combination of therapeutic agents. It can be administered alone, but will generally be administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

The dosage administered will vary depending on the use and known factors such as pharmacodynamic character of the particular agent, and its mode and route of administration; the recipient's age, weight, and health; nature and extent of symptoms; kind of concurrent treatment; frequency of treatment; and desired effect. For use in the treatment of said diseases or conditions, a compound of this invention can be orally administered



daily at a dosage of the active ingredient of 0.002 to 200 mg/kg of body weight. Ordinarily, a dose of 0.01 to 10 mg/kg in divided doses one to four times a day, or in sustained release formulation will be effective in obtaining the desired pharmacological effect.

Dosage forms (compositions) suitable for administration contain from about 1 mg to about 100 mg of active ingredient per unit. In these pharmaceutical compositions, the active ingredient will ordinarily be present in an amount of about 0.5 to 95% by weight based on the total weight of the composition.

The active ingredient can be administered orally in solid dosage forms, such as capsules, tablets and powders; or in liquid forms such as elixirs, syrups, and/or suspensions. The compounds of this invention can also be administered parenterally in sterile liquid dose formulations.

Gelatin capsules can be used to contain the active ingredient and a suitable carrier such as but not limited to lactose, starch, magnesium stearate, steric acid, or cellulose derivatives. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of time. Compressed tablets can be sugar-

coated or film-coated to mask any unpleasant taste, or used to protect the active ingredients from the atmosphere, or to allow selective disintegration of the tablet in the gastrointestinal tract.

- 5      Liquid dose forms for oral administration can contain coloring or flavoring agents to increase patient acceptance.

In general, water, pharmaceutically acceptable oils, saline, aqueous dextrose (glucose), and related  
10 sugar solutions and glycols, such as propylene glycol or polyethylene glycol, are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably contain a water soluble salt of the active ingredient, suitable stabilizing agents,  
15 and if necessary, buffer substances. Antioxidizing agents, such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or in combination, are suitable stabilizing agents. Also used are citric acid and its salts, and EDTA. In addition, parenteral  
20 solutions can contain preservatives such as benzalkonium chloride, methyl- or propyl-paraben, and chlorobutanol.

Suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences", A. Osol, a standard reference in the field.

Useful pharmaceutical dosage-forms for administration of the compounds of this invention can be illustrated as follows:

#### Capsules

5 A large number of units capsules are prepared by filling standard two-piece hard gelatin capsules each with 100 mg of powdered active ingredient, 150 mg lactose, 50 mg cellulose, and 6 mg magnesium stearate.

#### Soft Gelatin Capsules

10 A mixture of active ingredient in a digestible oil such as soybean, cottonseed oil, or olive oil is prepared and injected by means of a positive displacement was pumped into gelatin to form soft gelatin capsules containing 100 mg of the active  
15 ingredient. The capsules were washed and dried.

#### Tablets


A large number of tablets are prepared by conventional procedures so that the dosage unit was 100 mg active ingredient, 0.2 mg of colloidal silicon  
20 dioxide, 5 mg of magnesium stearate, 275 mg of microcrystalline cellulose, 11 mg of starch, and 98.8 mg lactose. Appropriate coatings may be applied to increase palatability or delayed adsorption.

The compounds of this invention may also be used as reagents or standards in the biochemical study of neurological function, dysfunction, and disease.

Although the present invention has been described  
5 and exemplified in terms of certain preferred  
embodiments, other embodiments will be apparent to those  
skilled in the art. The invention is, therefore, not  
limited to the particular embodiments described and  
exemplified, but is capable of modification or variation  
10 without departing from the spirit of the invention, the  
full scope of which is delineated by the appended  
claims.

What is claimed is:

1. A compound of Formula (I):



5

(I)

isomers thereof, stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, pharmaceutically acceptable prodrugs thereof, or  
10 pharmaceutically acceptable salt forms.

2. A compound of claim 1, pharmaceutically acceptable prodrugs thereof, or pharmaceutically acceptable salt forms thereof, wherein said compound is 4-((R)-2-  
15 butylamino)2,7-dimethyl-8-(2-methyl-6-methoxypyrid-3-yl)[1,5-a]-pyrazolo-1,3,5-triazine.

3. A compound of claim 1, pharmaceutically acceptable prodrugs thereof, or pharmaceutically acceptable salt  
20 forms thereof, wherein said compound is substantially free of its (S) stereoisomer.

4. A compound of claim 1, wherein said compound is 4-(2-butylamino)2,7-dimethyl-8-(2-methyl-6-methoxypyrid-3-yl) [1,5-a]-pyrazolo-1,3,5-triazine.

5. A compound of claim 1, wherein said compound is 4-((R)-2-butylamino)2,7-dimethyl-8-(2-methyl-6-methoxypyrid-3-yl) [1,5-a]-pyrazolo-1,3,5-triazine.

6. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of claim 1.

7. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of claim 5.

8. A method of antagonizing a CRF receptor in a mammal, comprising administering to the mammal, a therapeutically effective amount of a compound as claimed in claim 1.

9. A method of treating a disorder manifesting hypersecretion of CRF in a warm-blooded animal, comprising administering to the animal a therapeutically effective amount of a compound as claimed in claim 1.

10. A method for the treatment of a disorder, the treatment of which can be effected or facilitated by antagonizing CRF, comprising administering to the mammal a therapeutically effective amount of a compound of claim 1.

11. A method of antagonizing a CRF receptor in a mammal, comprising administering to the mammal, a therapeutically effective amount of a compound as claimed in claim 5.

12. A method of treating anxiety or depression in mammals, comprising administering to the mammal a therapeutically effective amount of a compound of claim 1.

13. A method of treating anxiety or depression in mammals, comprising administering to the mammal a therapeutically effective amount of a compound of claim 5.

14. A method for screening for ligands for CRF receptors, which method comprises:

- a) carrying out a competitive binding assay with a CRF receptor, a compound according to claim 1 which is labelled with a detectable label, and a candidate ligand; and
- b) determining the ability of said candidate ligand to displace said labelled compound.

30

15. A method for detecting CRF receptors in tissue comprising:
- a) contacting a compound of claim 1, which is labelled with a detectable label, with a tissue, under conditions that permit binding of the compound to the tissue; and
  - b) detecting the labelled compound bound to the tissue.
16. A method of inhibiting the binding of CRF to a CRF-1 receptor, comprising contacting a compound of claim 1 with a solution comprising cells expressing the CRF1 receptor, wherein the compound is present in the solution at a concentration sufficient to inhibit the binding of CRF to the CRF-1 receptor.
17. An article of manufacture comprising:
- a) a packaging material;
  - b) a compound of claim 1; and
  - c) a label or package insert contained within said packaging material indicating that said compound is effective for treating anxiety or depression.



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/06837

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) : A61P 25/00, 25/24; A61K 31/53; C07D 403/00

US CL : 514/245; 544/212

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/245; 544/212

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
CAS ONLINE**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 6,191,131 B1 (HE et al.) 20 February 2001, see entire document, especially Table 7.	1-17

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier application or patent published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\*

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\*

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\*

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*Z\*

document member of the same patent family

Date of the actual completion of the international search

03 June 2002 (03.06.2002)

Date of mailing of the international search report

11 JUL 2002

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks

Box PCT

Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Brenda L. Coleman

Telephone No. 703-308-1235